

NO-A191 903


LUF (LOWEST USABLE FREQUENCY) MODEL UNCERTAINTY
ASSESSMENT(U) NAVAL OCEAN SYSTEMS CENTER SAN DIEGO CA
D B SAILORS ET AL JAN 87

1/1

UNCLASSIFIED

F/G 20/14 NL





MICROCOPY RESOLUTION TEST CHART
 10-10-1963

AD-A191 903

UNCL/
SECURITY

REPORT DOCUMENTATION PAGE

1a REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b RESTRICTIVE MARKINGS	
2a SECURITY CLASSIFICATION AUTHORITY			3 DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution is unlimited.	
2b DECLASSIFICATION/DOWNGRADING SCHEDULE			5 MONITORING ORGANIZATION REPORT NUMBER(S)	
4 PERFORMING ORGANIZATION REPORT NUMBER(S)			7a NAME OF MONITORING ORGANIZATION	
6a NAME OF PERFORMING ORGANIZATION Naval Ocean Systems Center		6b OFFICE SYMBOL (if applicable) SPAWAR	7b ADDRESS (City, State and ZIP Code)	
8a NAME OF FUNDING/SPONSORING ORGANIZATION Space & Naval Warfare Systems Command			9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER	
8b ADDRESS (City, State and ZIP Code) Washington, DC 20363-5100			10 SOURCE OF FUNDING NUMBERS	
			PROGRAM ELEMENT NO RDDA	PROJECT NO MP57
			TASK NO NSA	AGENCY ACCESSION NO DN288 591
11 TITLE (Include Security Classification) LUF Model Uncertainty Assessment				
12 PERSONAL AUTHOR(S) D.B. Sailors				
13a TYPE OF REPORT Presentation		13b TIME COVERED FROM _____ TO _____		14 DATE OF REPORT (Year, Month, Day) January 1987
15 PAGE COUNT				
16 SUPPLEMENTARY NOTATION				
17 COSATI CODES			18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP		
19 ABSTRACT (Continue on reverse if necessary and identify by block number) <p>A statistical analysis of observed oblique sounder lowest observed frequency (LOF) and predicted lowest usable frequency (LUF) was used to assess the accuracy of the PROPHET algorithm QLOF version 2.0. A LOF data base of 1814 LOFs from 29 paths was established for the assessment. The data was screened into subsets to determine the effect of particular paths, universal and local time, sounder type, path length and orientation, season, month, sunspot number, and mid-path latitude. Version 2.0 had a bias of 0.55 MHz low, an rms error of 1.95 MHz, and a correlation coefficient with the observed data of 0.89. The rms error increases with increasing range. A higher residual was noted for the months of July through October. An increase in residual at mid-latitude was noted. Reasons for these errors are given. A modification to the absorption index term which determines the latitudinal and seasonal variation is suggested.</p>				
20 DISTRIBUTION/AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21 ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED	
22a NAME OF RESPONSIBLE INDIVIDUAL D.B. Sailors			22b TELEPHONE (Include Area Code) 619-225-7706	22c OFFICE SYMBOL Code 542

DD FORM 1473, 84 JAN


83 APR EDITION MAY BE USED UNTIL EXHAUSTED
ALL OTHER EDITIONS ARE OBSOLETEUNCLASSIFIED
SECURITY CLASSIFICATION OF THIS PAGEDTIC
ELECTE
MAR 11 1988

print

LUF MODEL UNCERTAINTY ASSESSMENT
D.B. Sailors and W.K. Moision
Ocean and Atmospheric Sciences Division
Naval Ocean Systems Center
San Diego, CA 92152-5000

A statistical analysis of observed oblique sounder lowest observed frequency (LOF) and predicted lowest usable frequency (LUF) was used to assess the accuracy of the PROPHET algorithm QLOF version 2.0. A LOF data base of 1814 LOFs from 29 paths was established for the assessment. The data was screened into subsets to determine the effect of particular paths, universal and local time, sounder type, path length and orientation, season, month, sunspot number, and mid-path latitude. Version 2.0 had a bias of 0.55 MHz low, an rms error of 1.95 MHz, and a correlation coefficient with the observed data of 0.89. The rms error increases with increasing range. A higher residual was noted for the months of July through October. An increase in residual at mid-latitude was noted. Reasons for these errors are given. A modification to the absorption index term which determines the latitudinal and seasonal variation is suggested.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Avail and/or	
Dist	Special
A-1	



END

DATE

FILMED

5-88

DTIC